

Hulk

PEPTIDES

THE HULK: 10MG BLEND – CYCLE, ADMINISTRATION, GENERAL CONSIDERATIONS 8-12 week cycle followed by 6-8 week washout period If you are doing an 8 week cycle, dose 7 days per week; if doing a 12-week cycle, dose 5 days per week Dose once per day Bedtime dosing recommended; may also b

THE HULK: 10MG BLEND – CYCLE, ADMINISTRATION, GENERAL CONSIDERATIONS 8-12 week cycle followed by 6-8 week washout period If you are doing an 8 week cycle, dose 7 days per week; if doing a 12-week cycle, dose 5 days per week Dose once per day Bedtime dosing recommended; may also be dosed immediately upon waking in the a.m Subcutaneous administration To avoid hot/flushed/prickly skin sensation, inject shot as slowly as possible Reconstitution Mix with 2.5mL (250) units of BAC water Dosing Weeks 1-2: 3 units (100mcg), 1x/day, 5-7 days per week Weeks 3-4: 4 units (150mcg), 1x/day, 5-7 days per week Weeks 5-6: 5 units (200mcg), 1x/day, 5-7 days per week Weeks 7-8: 6 units (250mcg), 1x/day, 5-7 days per week Weeks 9-10: 8units (300mcg), 1x/day, 5-7 days per week Weeks 11-12: 9 units (350mcg), 1x/day, 5-7 days per week

RESEARCH USE DISCLAIMER: All information in this document is provided strictly for laboratory research and educational purposes. It is not intended for human or veterinary use, diagnosis, or treatment.