

KPV

PEPTIDES

KPV is a C-terminal tripeptide of α -MSH (alpha-Melanocyte-stimulating hormone) that acts in a manner similar to hormones and relays information from one tissue through the blood to another via biological messengers. Helpful in the treatment of a broad array of conditions, from skin disorders to

KPV is a C-terminal tripeptide of α -MSH (alpha-Melanocyte-stimulating hormone) that acts in a manner similar to hormones and relays information from one tissue through the blood to another via biological messengers. Helpful in the treatment of a broad array of conditions, from skin disorders to musculoskeletal conditions to gut problems, and many more. Excellent anti-inflammatory benefits, making it useful in treating various inflammatory conditions Boosts the immune system Promotes wound healing and faster recovery of damaged tissues Improves skin health (including conditions like acne and psoriasis) Enhances gut health (helpful for reducing symptoms of Inflammatory Bowel Disease) Has antimicrobial effects against pathogens such as *S. aureus* and *C. albicans* Aids recovery from liver injury by increasing the replication of liver cells Reduces scarring and improves the organization of collagen fibers Improves the prolonged survival of grafted tissues Suppresses inflammatory responses which spur contact dermatitis Can help fight fungal infections POSSIBLE SIDE EFFECTS Injection site reactions (redness, swelling, itching) Nausea Fatigue CONTRAINDICATIONS Do not take if you are pregnant or breastfeeding Do not take if you have an active cancer diagnosis RECONSTITUTION, DOSING , & OTHER CONSIDERATIONS 8-week cycle followed by 4-8 week washout period If you are using KPV for gut-repair, dose any time day or night, but designate a specific time each day that you will be able to dose: consistent dose time is important for this purpose If you are using KPV for anything other than gut repair, dose any time of day Daily dosing Mix with 2.5mL (250 units) of BAC water 8 week cycle followed by 4-6 week washout period First dose is 5 units (200 mcg), once per day Increase dose in 3 unit (100mcg) intervals, only as necessary to continue sustaining benefits Maximum dose is 25 units (1mg) 100mcg is actually 2.5 units, but since it is absolutely impossible to accurately measure this on any syringe, we have rounded up to 3 units.

RESEARCH USE DISCLAIMER: All information in this document is provided strictly for laboratory research and educational purposes. It is not intended for human or veterinary use, diagnosis, or treatment.