

NAD-Glutathione-Vitamin-C-Intranasal

PEPTIDES

CONTRAINDICATIONS Do not use if pregnant or nursing Do not use if you have had a recent or active cancer diagnosis Do not use if you are hypersensitive or allergic to Glutathione or NAD+ **DOSING** Cycle Length: 12 weeks followed by 4 week washout period. Dosing Times: Morning prefer

CONTRAINDICATIONS Do not use if pregnant or nursing Do not use if you have had a recent or active cancer diagnosis Do not use if you are hypersensitive or allergic to Glutathione or NAD+ **DOSING** Cycle Length: 12 weeks followed by 4 week washout period. Dosing Times: Morning preferred, as NAD+ may lead to increased energy and sleep disturbance. This possibility should be taken into consideration by those who wish to dose a second time per day. Onset Time: 15 minutes to 1 hour. Additional Considerations: Does not require administration during a fasted state, as the intranasal administration bypasses the gastrointestinal tract. Storage: Atomizer should be refrigerated. Administration: Intranasal spray. Always split dose between both nostrils as much as possible. This is because it increases the mucosal surface area exposed to the solution, decreases runoff into the throat/loss from swallowing, and allows for more consistent absorption because nasal airflow, congestion, mucus, and the normal nasal cycle may make one nostril less effective than the other at any given time. Proper technique is important. **DO NOT TILT YOUR HEAD BACK** to dose. With your head tipped forward , tilt your atomizer towards the outside of your nose/outside corner of your eye, then sniff the spray gently. 2-4 sprays, once or twice per day, daily.

RESEARCH USE DISCLAIMER: All information in this document is provided strictly for laboratory research and educational purposes. It is not intended for human or veterinary use, diagnosis, or treatment.