

Orforglipron-2

PEPTIDES

Orforglipron is glucagon-like peptide-1 (GLP-1RA) weight that increases the release of insulin from the pancreas by increasing the volume of insulin-producing beta cells and reducing the release of glucagon (a hormone that increases blood sugar levels). Primarily a weight loss dr

Orforglipron is glucagon-like peptide-1 (GLP-1RA) weight that increases the release of insulin from the pancreas by increasing the volume of insulin-producing beta cells and reducing the release of glucagon (a hormone that increases blood sugar levels). Primarily a weight loss drug, it works by slowing the movement of food from the stomach into the small intestine, resulting in increased satiety, reduced food intake, and more efficient energy expenditure. DOSING CONSIDERATIONS 18-36 week cycle followed by 18-36 week washout period May be taken at any time of day Does not have to be taken while in a fasted state There are no dosing-related water intake restrictions, as seen in previous oral GLP formulation DOSING To dose accurately, please purchase a pill splitter. We have included links to appropriate options below: [PILL SPLITTER #1](#) [PILL SPLITTER #2](#) Week 1: 1/4 of a pill (1.5mg), once per day Week 2: 1/2 of a pill (3mg), once per day Weeks 3-6: 1 pill (6mg), once per day Weeks 7-10: 2 pills (12mg), once per day Weeks 11-14: 3 pills (18mg), once per day Weeks 15-18: 4 pills (24mg), once per day Weeks 19-22: 5 pills (30mg), once per day Weeks 24-36: 6 pills (36mg), once per day After 36 weeks, take a break for 18-36 weeks before starting new cycle

RESEARCH USE DISCLAIMER: All information in this document is provided strictly for laboratory research and educational purposes. It is not intended for human or veterinary use, diagnosis, or treatment.