

PT-141

PEPTIDES

IMPORTANT: Read This Before Proceeding With Dosing Go to the Prep & Injection Guide for proper reconstitution, syringe sizing, and injection protocols. Mistakes here can compromise your research. the Prep & Injection Guide PT-141 PT-141 (aka bremelanotide) is

IMPORTANT: Read This Before Proceeding With Dosing Go to the Prep & Injection Guide for proper reconstitution, syringe sizing, and injection protocols. Mistakes here can compromise your research. the Prep & Injection Guide PT-141 PT-141 (aka bremelanotide) is a synthetic peptide for treating sexual dysfunction and other related issues. It is intended to enhance sexual desire and increase sexual satisfaction in both men and women. It works by stimulating the melanocortin system, which is responsible for regulating sexual arousal, among other functions. Not only is there a remarkable enhancement of libido, but increased orgasmic responses as well. Benefits Include Reduced sexual anxiety and improved confidence Non-invasive treatment option with fewer side effects than alternatives like Viagra Long-lasting effects compared to other treatments Treats hypoactive sexual desire disorder in premenopausal women Improves erectile quality in some men Heightened genital sensitivity Reduced performance anxiety secondary to improved arousal Improved sexual confidence Does not have adverse reactions with alcohol Common Side Effects Flushing Headaches Nausea Irritation Injection site irritation (redness, swelling, and/or itching) Headaches and flushing are the most common side effects, reported in about 20–40% of users. Nausea occurs in about 15% of users. Regardless, these side effects are generally mild in severity. Less Common but More Serious Side Effects Include Vomiting Blurred vision Dizziness Joint pain or swelling Runny or stuffy nose Fatigue Please note: Dizziness and blurred vision occurring simultaneously may indicate a more serious underlying issue. Consult a physician immediately if this happens. Dosing Protocol PT-141: 10 mg vial Mix with 2 mL (200 units) of bacteriostatic water Subcutaneous Administration Inject 45 minutes to 1 hour prior to sexual activity Initial Dose (Test for Tolerance) 10 units (500 mcg) This is not intended to be effective for sexual activity. It ensures your body tolerates the peptide before using a full therapeutic dose. Use this dose before an actual encounter. You may need to build up to 40 units if you experience adverse side effects from the initial dose. Standard Dose 40 units (2 mg) Inject 45–60 minutes before sex Do not exceed 1 injection in 24 hours Do not exceed 8 standard doses (16 mg total) per month Maximum Dose 50 units (2.5 mg) Only use this dose if you've developed a tolerance to the standard dose Do not exceed this dose under any circumstances Do not exceed 8 total doses per month Important Notes Long-term use of PT-141 may lead to desensitization of the melanocortin system, reducing effectiveness If you need the effects of PT-141 more frequently, ask about alternate peptides to rotate between for safety DO NOT exceed monthly maximums — this could damage your health PT-141 Stacking Suggestions Melanotan-2 – for enhanced libido, skin tone, and sexual confidence Oxytocin – amplifies connection, bonding, and emotional intimacy effects CJC-1295 / Ipamorelin – improve recovery, energy, and general hormone optimization, complementing libido benefits BPC-157 /

TB-500 – for overall tissue recovery and performance support, especially in physically active users
Vitamin B12 – supports energy and can reduce nausea side effects commonly experienced with
PT-141 For safe long-term use, rotate PT-141 with synergistic or alternate sexual enhancement
peptides instead of overusing it.

RESEARCH USE DISCLAIMER: All information in this document is provided strictly for laboratory research and educational purposes. It is not intended for human or veterinary use, diagnosis, or treatment.