

Semax-Intranasal

PEPTIDES

DOSING Cycle Length: 4-6 weeks followed by 2-4 week washout period Dosing Times: Morning to early evening is ideal. 1-3 doses per day. Onset Time: 15-30 minutes. Additional Considerations: Does not require administration during a fasted state, as the intranasal administration bypasses the

DOSING Cycle Length: 4-6 weeks followed by 2-4 week washout period Dosing Times: Morning to early evening is ideal. 1-3 doses per day. Onset Time: 15-30 minutes. Additional Considerations: Does not require administration during a fasted state, as the intranasal administration bypasses the gastrointestinal tract. Storage: Semax atomizer should be refrigerated after use. Administration: Intranasal spray. Always split dose between both nostrils as much as possible. This is because it increases the mucosal surface area exposed to the solution, decreases runoff into the throat/loss from swallowing, and allows for more consistent absorption because nasal airflow, congestion, mucus, and the normal nasal cycle may make one nostril less effective than the other at any given time. Dispense Rate: 100mcg per spray Standard Protocol: 2-5 sprays (200-500mcg), 1-3 times per day Maximum Protocol: 5-10 sprays (500mcg- 1mg), 1- 3 times per day *Dose listed is not divided between 1-3 times per day, but is to be taken each time you dose.

RESEARCH USE DISCLAIMER: All information in this document is provided strictly for laboratory research and educational purposes. It is not intended for human or veterinary use, diagnosis, or treatment.