

Slu-Pp-332-Oral

PEPTIDES

CYCLE LENGTH& WASHOUT PERIOD: 4-8 week cycle followed by 2-4 week washout period.

DOSING FREQUENCY: Because of the short cycle length, daily dosing will not lead to desensitization, and is therefore recommended. However, 5 consecutive dosing days followed by 2 rest days is also a

CYCLE LENGTH& WASHOUT PERIOD: 4-8 week cycle followed by 2-4 week washout period.

DOSING FREQUENCY: Because of the short cycle length, daily dosing will not lead to desensitization, and is therefore recommended. However, 5 consecutive dosing days followed by 2 rest days is also acceptable. TIME OF DAY: Morning. FASTED?: Yes, you should be in a fasted state to dose.

ADMINISTRATION: Oral (capsules). SLU-PP-332, CAPSULES: 300MCG Week 1: Take one capsule (300mcg) in the morning, daily. Weeks 2 through End of Cycle: Take 1-2 capsules (300-600mcg), once per day. SLU-PP-332, CAPSULES: 1,000MCG Take one capsule (1,000mcg) in the morning, daily, for the duration of the cycle. NOTE: The 1,000mcg capsules are not intended for first-time SLU-PP-332 users. Should be limited to those who are on their second or later cycle, and who have tolerated lower doses without substantial adverse side effects.

RESEARCH USE DISCLAIMER: All information in this document is provided strictly for laboratory research and educational purposes. It is not intended for human or veterinary use, diagnosis, or treatment.