

# Slupporforglipron

## PEPTIDES

POSSIBLE SIDE EFFECTS Nausea Constipation Diarrhea Heartburn Fatigue (particularly during dose increases) Increased body temperature Increased sweating Mild insomnia DOSING: SLU-PP-332 (250mcg) + ORFORGLIPRON (6mg) Take in the morning Does not have to be taken while fasted, but t

POSSIBLE SIDE EFFECTS Nausea Constipation Diarrhea Heartburn Fatigue (particularly during dose increases) Increased body temperature Increased sweating Mild insomnia DOSING: SLU-PP-332 (250mcg) + ORFORGLIPRON (6mg) Take in the morning Does not have to be taken while fasted, but taking it while fasted will lead to better fat loss Maintain a high-protein diet and sufficient hydration Week 1: Take 1 pill every other day (days 1, 3, 5, and 7) if you have never taken any form of GLP before. For those who have taken any form of GLP at any point in time, skip Week 1 and go immediately to Week 2. Week 2 – 10: Take 1 pill every day (so long as you are not experiencing substantial adverse side effects) Week 11+: Take 2 pills every day

*RESEARCH USE DISCLAIMER: All information in this document is provided strictly for laboratory research and educational purposes. It is not intended for human or veterinary use, diagnosis, or treatment.*