

Wolverineblend

PEPTIDES

RECONSTITUTION AND DOSING PROTOCOLS FOR BPC-157+TB-500: 10MG "WOLVERINE" BLEND 8-week cycle followed by 4-8 week washout period May be taken at any time of day Dosing does not require fasting Subcutaneous or Intramuscular Administration RECONSTITUTION Mix with 2mL (200 units) of

RECONSTITUTION AND DOSING PROTOCOLS FOR BPC-157+TB-500: 10MG "WOLVERINE" BLEND 8-week cycle followed by 4-8 week washout period May be taken at any time of day Dosing does not require fasting Subcutaneous or Intramuscular Administration RECONSTITUTION Mix with 2mL (200 units) of BAC water DOSING STANDARD PROTOCOL Weeks 1-2: 6 units (600mcg), 1x/day Weeks 3-8: 8 units (800mcg), 1x/day MAXIMUM PROTOCOL Weeks 1-2: 10 units (1,000mcg), 1x/day Weeks 3-8: 6 units (600mcg), 1x/day ¹: Either subcutaneous or intramuscular injection may be used to administer this blend. Each has it's own pro's and con's. You may rotate between the two at will. However, certain regions of the body result in higher systemic bioavailability, with the stomach offering the best absorption overall. Subcutaneous: Pro's: Provides systemic benefits by entering the bloodstream and circulating throughout the body; good for both general healing and healing multiple injuries. Con's: Absorption is slower than in IM injections. Intramuscular: Pro's: Faster absorption and more noticeable effects. Suitable for acute injuries, post-surgery recovery, or severe inflammation where quick effects are needed. Can be used for general benefits and systemic benefits and is also recommended for targeted pain relief, especially localized injections near the affected area. Con's: Some may consider IM injections to be more difficult and/or more painful.

RESEARCH USE DISCLAIMER: All information in this document is provided strictly for laboratory research and educational purposes. It is not intended for human or veterinary use, diagnosis, or treatment.